

Zero sets of entire functions of exponential type with some conditions on real axis

Favorov S.Ju.

Abstract

We give a complete description of zero sets for some well-known subclasses of entire functions of exponential growth.

2000 Mathematics Subject Classification: Primary 30D15, Secondary 30D10, 30D45

Keywords: entire function of exponential type, Cartwright's function, zero set of entire function

By B we denote the class of entire functions of exponential type bounded on the real axis;

by C denote the class of Cartwright's entire functions, i.e., entire functions f of exponential type with the property

$$\int_{-\infty}^{\infty} \frac{\log^+ |f(x)|}{1+x^2} dx < \infty, \quad (1)$$

by D denote the class of entire functions f of exponential type with the property: for each $h_m \in \mathbb{R}$ there exists a subsequence h'_m and a function $\tilde{f} \not\equiv 0$ such that the sequence $f(z + h'_m)$ converges uniformly to a function $\tilde{f}$ on every compact subset of $\mathbb{C}$.

It is easy to see that $D \subset B \subset C$. The classes A and B are well known and prove useful in various areas of complex and functional analysis (cf. [1], [2], [3]), class D is an extension of the well known class of Sine-Type functions (cf.[2]); it inherits some properties of Sine-Type functions.

Here we will consider sequences $\{a_k\}$ of complex numbers without finite limit points; note that each $\{a_k\}$ may appear in the sequence with a finite multiplicity. We say that a sequence $\{a_k\}$ is a *zero set for an entire function* f if $f(a) \neq 0$ for $a \notin \{a_k\}$ and $f(a) = f'(a) = \dots = f^{(p-1)}(a) = 0$, $f^{(p)}(a) \neq 0$, where p is a multiplicity of a in $\{a_k\}$.

The zero set of any function $f \in C$ satisfies the conditions

$$\sum_{a_k \neq 0} |\Im a_k^{-1}| < \infty, \quad (2)$$

$$\exists \lim_{R \rightarrow \infty} \sum_{0 < |a_k| < R} a_k^{-1}, \quad (3)$$

and for all $0 < \alpha \leq \pi/2$

$$\lim_{R \rightarrow \infty} R^{-1} \text{card}\{a_k : 0 < |a_k| < R, |\arg a_k| \leq \alpha\} = d/2\pi, \quad (4)$$

$$\lim_{R \rightarrow \infty} R^{-1} \text{card}\{a_k : 0 < |a_k| < R, |\arg a_k - \pi| \leq \alpha\} = d/2\pi, \quad (5)$$

where d is the width of the indicator diagram of f (cf.[2], p.127). Then every function $f \in C$ has the form

$$f(z) = Az^s e^{i\lambda z} \lim_{R \rightarrow \infty} \prod_{0 < |a_k| < R} (1 - z/a_k) \quad (6)$$

with $A \in \mathbb{C}$, $\lambda \in \mathbb{R}$ (cf.[2], p.130). However there are zero sets satisfying (2)–(5) such that every entire function with this zero set does not belong to this class ¹.

B. Levin in [1], Appendix VI, investigated zero sets $\{a_k\}$, $k \in \mathbb{Z}$, for functions of class B with the additional property $\sup_k |a_k - hk| < \infty$ with some

¹For example, take $a_1 = -e^2$ and reals $a_k < a_1$, $k = 2, \dots$, such that the number $n(r) = \text{card}\{a_k : a_k \geq -r\}$ is the integer part of $r/\log^2 r$ for each $r \geq e^2$. Then (2)–(5) fulfilled and the function (6) with $s = 0$ does not satisfy (1) because for $x > e^2$ we have

$$\log |f(x)| = \int_1^\infty \log(1 + x/t) dn(t) = \int_1^\infty \frac{xn(t)dt}{t(x+t)} > 1 + \int_x^\infty \frac{xdt}{2t \log^2 t} = 1 + \frac{x}{2 \log x};$$

the idea belongs to prof. A. Grishin.

$h \in \mathbb{R}$. In the present paper we obtain a complete description of zero sets for the classes B , C , D .

In what follows we put $n(c, t) = \text{card}\{a_k : |a_k - c| \leq t\}$ for $c \in \mathbb{C}$. Assume also that the sequence $\{a_k\}$ satisfies the following conditions

$$n(0, t) = O(t), \quad t \rightarrow \infty, \quad (7)$$

and

$$n(0, t+1) - n(0, t) = o(t), \quad t \rightarrow \infty. \quad (8)$$

It is clear that (7) and (8) are invariant with respect to the change $n(0, t)$ to $n(c, t)$.

Theorem 1 *A sequence $\{a_k\} \subset \mathbb{C}$ is the zero set of some function $f \in C$ if and only if it satisfies (3), (7), (8), and*

$$\int_{-\infty}^{\infty} \left[\int_0^{\infty} [n(b, t) - n(x, t)] t^{-1} dt \right]^+ \frac{dx}{1+x^2} < \infty, \quad (9)$$

for some point $b \in \mathbb{R} \setminus \{a_k\}$.

Theorem 2 *A sequence $\{a_k\} \subset \mathbb{C}$ is the zero set of some function $f \in B$ if and only if it satisfies (3), (7), (8), and*

$$\sup_{x \in \mathbb{R}} \int_0^{\infty} [n(b, t) - n(x, t)] t^{-1} dt < \infty \quad (10)$$

for some point $b \in \mathbb{R} \setminus \{a_k\}$ ².

Theorem 3 *A sequence $\{a_k\} \subset \mathbb{C}$ is the zero set of some function $f \in D$ if and only if it satisfies (3), (7), (8), and*

$$\sup_{x \in \mathbb{R}} \left| \int_1^{\infty} [n(0, t) - n(x, t)] t^{-1} dt \right| < \infty. \quad (11)$$

The proofs of Theorems 1–3 are based on the following lemma:

²Theorems 1 and 2 were announced (with my permission) in survey [4], p.45.

Lemma 1 . Let a sequence $\{a_k\} \subset \mathbb{C} \setminus \{0\}$ satisfy (3), (7), and (8). Then

$$g(z) = \lim_{R \rightarrow \infty} \prod_{|a_k| < R} (1 - z/a_k) \quad (12)$$

is a well-defined entire function of a finite exponential type, and for all $z \in \mathbb{C}$

$$\log |g(z)| = \int_0^\infty [n(0, t) - n(z, t)] t^{-1} dt. \quad (13)$$

Moreover, if z_0 is a point from the sequence $\{a_k\}$ with multiplicity $l = n(z_0, 0) > 0$, then

$$\log \left| \frac{g^{(l)}(z_0)}{l!} \right| = \int_1^\infty \frac{n(0, t) - n(z_0, t)}{t} dt + \int_0^1 \frac{n(0, t) - n(z_0, t) + l}{t} dt. \quad (14)$$

Proof of Lemma. It follows from (7) that the integral $\int_0^\infty n(0, t) t^{-3}$ is finite, and so

$$\sum |a_k|^{-2} = \int_0^\infty t^{-2} dn(0, t) < \infty. \quad (15)$$

Let K be a fixed disc in the complex plane. Then

$$\sum_{R < |a_k| < R'} \log \left(1 - \frac{z}{a_k} \right) - \sum_{R < |a_k| < R'} \frac{z}{a_k} \rightarrow 0$$

as $R, R' \rightarrow \infty$ uniformly on $z \in K$. Hence, uniformly on $z \in K$ we have

$$\lim_{R \rightarrow \infty} \sum_{|a_k| > R} \log \left(1 - \frac{z}{a_k} \right) = 0, \quad (16)$$

and the function g is well-defined. Conditions (3) and (7) now yield that g (12) has a finite exponential type of growth.

Next, for $z = a_k$ equality (13) is trivial. Fix $z \notin \{a_k\}$. Take $\varepsilon > 0$ such that $n(0, \varepsilon) = n(z, \varepsilon) = 0$. Then we have for any $R < \infty$

$$\begin{aligned} \sum_{|a_k| \leq R} (\log |z - a_k| - \log |a_k|) &= \sum_{\varepsilon \leq |a_k - z| \leq R} \log |z - a_k| - \sum_{\varepsilon \leq |a_k| \leq R} \log |a_k| \\ &+ \sum_{|a_k| \leq R, |z - a_k| > R} \log |z - a_k| - \sum_{|z - a_k| \leq R, |a_k| > R} \log |z - a_k|. \end{aligned} \quad (17)$$

Furthermore,

$$\sum_{\varepsilon \leq |a_k - z| \leq R} \log |z - a_k| = n(z, R) \log R - \int_{\varepsilon}^R n(z, t) t^{-1} dt,$$

and

$$\begin{aligned} & (\log R)[n(z, R) - n(0, R)] + \sum_{|a_k| \leq R, |z - a_k| > R} \log |z - a_k| - \sum_{|z - a_k| \leq R, |a_k| > R} \log |z - a_k| \\ &= \sum_{|a_k| \leq R, |z - a_k| > R} \log \frac{|z - a_k|}{R} + \sum_{|z - a_k| \leq R, |a_k| > R} \log \frac{R}{|z - a_k|}. \end{aligned} \quad (18)$$

Note that if $|z - a_k| > R$ and $|a_k| \leq R$, then $|a_k| > R - |z|$ and $|z - a_k| \leq |z| + R$. Also, if $|a_k| > R$ and $|z - a_k| \leq R$, then $|a_k| \leq R + |z|$ and $|z - a_k| > R - |z|$. Hence for $R \rightarrow \infty$ the right-hand side of (18) does not exceed

$$O(|z|/R)[n(0, R) - n(0, R - |z|)] + O(|z|/R)[n(0, R + |z|) - n(0, R)]. \quad (19)$$

Taking into account (8), we see that (19) tends to zero for fixed z as $R \rightarrow \infty$. Therefore assertion (13) follows from (17) and (16). In order to get (14), apply (13) to the function $g(z)(z - z_0)^{-l} z_0^l$. Lemma is proved.

Proof of Theorems 1–3. It follows from (15) that the limits as $R \rightarrow \infty$ of the sums $\sum_{|a_k - c| < R} (a_k - c)^{-1}$ and $\sum_{|a_k - c| < R} (a_k)^{-1}$ exist simultaneously, and the last sum differs from the sum $\sum_{|a_k| < R} (a_k)^{-1}$ no more than $n(0, R + |c|) - n(0, R - |c|) = o(R)$ terms, where modulus of each term does not exceed $(R - |c|)^{-1}$. Therefore (3) is invariant with respect to the shift of the origin. Further, using Lemma 1 for the function $\lim_{R \rightarrow \infty} \prod_{|a_k - c| < R} (1 - z/(a_k - c))$, we see that $[\int_0^\infty [n(c, t) - n(x, t)] t^{-1} dt]^+ = O(|x|)$ as $|x| \rightarrow \infty$, therefore (9) is invariant with respect to the shift on a real b as well. Hence with no loss of generality we can assume $0 \notin \{a_k\}$ and $b = 0$ in (9), (10).

It follows immediately from Lemma 1 that under conditions of Theorems 1 or 2 the function (12) belongs to class C or B , respectively. Let $\{a_k\}$ satisfy the conditions of Theorem 3. By (13), (14), and Jensen formula

$$\frac{1}{2\pi} \int_0^{2\pi} \log |g(z + e^{i\theta})| d\theta = \log \left| \frac{g^{(l)}(z)}{l!} \right| + \int_0^1 \frac{n(z, t) - n(z, 0)}{t} dt$$

with $l = n(z, 0)$, we get

$$\frac{1}{2\pi} \int_0^{2\pi} \log |g(z + e^{i\theta})| d\theta = \int_1^\infty \frac{n(0, t) - n(z, t)}{t} dt + \int_0^1 \frac{n(0, t)}{t} dt. \quad (20)$$

Hence it follows from (11) and the inequality

$$\log |g(z)| \leq \frac{1}{2\pi} \int_0^{2\pi} \log |g(z + e^{i\theta})| d\theta,$$

that g is bounded on $\mathbb{R}$. The Pfragmen–Lindelof Theorem implies that $g(z)$ is bounded on every horizontal strip of a finite width. Therefore for each $t_m \in \mathbb{R}$ there exists a subsequence t'_m such that $f(z + t'_m)$ converge uniformly to a function $h(z)$ on every compact subset of $\mathbb{C}$. If $h(z) \equiv 0$ on $\mathbb{C}$, then $\int_0^{2\pi} \log |g(t'_m + e^{i\theta})| d\theta \rightarrow -\infty$. By (20), this contradicts (11), and so $g \in D$.

To prove necessity, note that by (6) every function $f \in C$ with $f(0) \neq 0$ has the form

$$f(z) = Ae^{i\lambda z} g(z)$$

with g defined in (12). The conditions (7) and (8) follow immediately from (4) and (5). Using Lemma 1, we see that the zero set of f satisfies (9) and, when $f \in B$, (10). For the case $f \in D$ the functions f and g are bounded on $\mathbb{R}$ and, by the Pfragmen–Lindelof Theorem, on every horizontal strip of a finite width. Hence, if (11) is false, then from (20)

$$\frac{1}{2\pi} \int_0^{2\pi} \log |g(t_l + e^{i\theta})| d\theta \rightarrow -\infty$$

for some sequence of reals t_j . Taking into account properties of averages of subharmonic functions, we have

$$\int_{|u+iv-t_j|<1} \log |g(u+iv)| du dv \rightarrow -\infty.$$

Since the function $\log |g|$ is bounded from above on the strip $|\Im z| < 2$, we get

$$\int_{|u+iv-t_j-s|<2} \log |g(u+iv)| du dv \rightarrow -\infty$$

uniformly in $s \in [-1, 1]$. Using properties of averages of subharmonic functions again, we obtain

$$\sup_{-1 \leq s \leq 1} \log |g(t_j + s)| \rightarrow -\infty. \quad (21)$$

On the other hand, $f(z+t'_j) \rightarrow h(z) \not\equiv 0$ for some subsequence t'_j uniformly on compact subsets of $\mathbb{C}$, hence (21) is impossible. Theorems 1–3 are completely proved.

Remark 1. It can be shown that all these theorems are valid with the function $\tilde{n}(c, t) = \text{card}\{a_k : |\Re(a_k - c)| \leq t, |\Im(a_k - c)| \leq t\}$ instead of $n(c, t)$ as well.

Remark 2. The additional condition

$$\limsup_{y \rightarrow \pm\infty} \int_0^\infty [n(b, t) - n(iy, t)](|y|t)^{-1} dt \leq \sigma$$

in Theorems 1–3 gives the complete description for zero sets of corresponding subclasses of functions with an exponential type at most σ .

Example. Let $\alpha(t)$ be a strictly increasing concave function on $\mathbb{R}^+$ with the properties $\alpha(0) = 0$, $1 \leq \alpha'(t) \leq 1 + O(t^{-1})$ as $t \rightarrow \infty$, $\alpha(t) \geq t + 1 + \varepsilon$ for large t with some $\varepsilon > 0$. Consider a sequence of reals $\{a_k\}$, $k \in \mathbb{Z} \setminus \{0\}$, such that $\alpha(a_k) = k$ and $\{a_{-k}\} = \{-a_k\}$, $k \in \mathbb{N}$. Note that

$$n(0, t) - n(x, t) = 2E[\alpha(t)] + E[\alpha(x - t)] - E[\alpha(x + t)]$$

for $x > t$, and

$$n(0, t) - n(x, t) = 2E[\alpha(t)] - E[\alpha(x - t)] - E[\alpha(x + t)]$$

for $0 < x < t$; here $E[x]$ is the integer part of x . Since $a_{n+1} - a_n = (\alpha'(\tilde{t}))^{-1}$ with some $\tilde{t} \in (a_n, a_{n+1})$, we see that for $n \geq n_0$

$$0 \leq 1 - (a_{n+1} - a_n) \leq (a_{n+1} - a_n)^{-1} - 1 \leq \alpha'(a_n) - 1 < C/a_n < 1/2,$$

and

$$\left| \int_{a_n}^{a_{n+1}} (\alpha(t) - E[\alpha(t)] - 1/2) dt \right| \leq \left| \int_{a_n}^{a_{n+1}} \left(\int_{a_n}^t \alpha'(u) du - 1/2 \right) dt \right| \leq C'/a_n.$$

Therefore we have for arbitrary $N < \infty$

$$\left| \int_{a_{n_0}}^{a_N} (\alpha(t) - E[\alpha(t)] - 1/2) dt \right| \leq C' \log a_N.$$

Now it follows easily that all the integrals

$$\int_1^x \frac{\alpha(t) - E[\alpha(t)] - 1/2}{t} dt, \quad \int_1^x \frac{\alpha(x \pm t) - E[\alpha(x \pm t)] - 1/2}{t} dt,$$

$$\int_x^\infty \frac{\alpha(t \pm x) - E[\alpha(t \pm x)] - 1/2}{t} dt,$$

are bounded uniformly in $x \rightarrow \infty$. Hence, up to some uniformly bounded term, integral $\int_0^\infty [n(0, t) - n(x, t)] t^{-1} dt$ equals

$$\begin{aligned} \int_1^x \frac{2E[\alpha(t)] - 2t}{t} dt + \int_1^{x-1} \frac{\alpha(x-t) - \alpha(x+t) + 2t}{t} dt \\ + \int_x^\infty \frac{2\alpha(t) - \alpha(t-x) - \alpha(x+t)}{t} dt. \end{aligned} \quad (22)$$

The function $\alpha(t)$ is concave, therefore the last integral in (22) is positive. Further, we have for $1 < t < x-1$

$$\alpha(x-t) - \alpha(x+t) + 2t = \int_{-t}^t 1 - \alpha'(x+u) du \geq -C \log \frac{x+t}{x-t}.$$

Now it is easy to check that the second integral in (22) is uniformly bounded from below. Evidently, the first integral in (22) is unbounded from above as $x \rightarrow \infty$. Thus the sequence $\{a_k\}$ is the zero set of no function $f \in B$.

References

- [1] Levin, B.Ja., Distributions of Zeros of Entire Functions. Transl. of Math. Monograph, Vol.5, AMS Providence, R1, 1980.
- [2] Levin, B.Ja., Lectures on Entire Functions. Transl. of Math. Monograph, Vol.150, AMS Providence, 1996, 248p.
- [3] Koosis, P., The logarithmic integral. I, Cambridge Univ. Press, Cambridge, 1988, II, Cambridge Univ. Press, Cambridge, 1992.
- [4] Khabibullin, B., Completeness of Exponent Sistems and Sets of Uniqueness. Rits BashGU, Ufa, 2006, (russian).

Department of Mathematics, Kharkov National University,
Svobody sq.,4, Kharkov 61077, Ukraine